Reading:

• Hughes and Hase, Chapter 4

Problems due Tuesday February 3:

1. In Table 4.2 on p. 44, Hughes & Hase claim that for the function

$$Z(A,B) = k \frac{A^n}{B^m}$$

the fractional uncertainty in Z is given by

$$\frac{\alpha_Z}{Z} = \sqrt{\left(n\frac{\alpha_A}{A}\right)^2 + \left(m\frac{\alpha_B}{B}\right)^2}.$$

Use the *calculus approach* to prove this result.

2. In the PHYS 211/212 appendix we discuss the uncertainty in the measurement of g with a pendulum. We say that the uncertainty in the value of g due to our uncertainty in the measurement of the period T is given by

$$\Delta g_T = g(L, T + \Delta T) - g(L, T)$$

(This is H&H's functional approach. It's just as reasonable to use

$$\Delta q_T = q(L, T - \Delta T) - q(L, T).$$

Let's assume that L = 0.96 m and $T = 1.970 \pm 0.004$ s. Does it matter which definition for Δg_T you use? For what values of ΔT will it matter?

- 3. Section 4.2.2 in Hughes & Hase is a worked example of the determination of pressure and its uncertainty using the van der Waals equation of state and the functional approach for determining uncertainties. Repeat these calculations for yourself, determining $P(\bar{V}_{in}, \bar{T})$, α_P^T , α_P^V , and α_P . Identify the (slight) numerical errors made in the text.
- 4. Repeat the calculation of the uncertainty α_P in problem #3 using the "calculus approximation" of the uncertainties.

- 5. Repeat the calculation of the uncertainty α_P in problem # 3 using Monte Carlo simulations of the data.
- 6. Hughes and Hase, Problem 4.8
- 7. Hughes and Hase, Problem 4.10