Homework 10

Date Assigned: Thursday, April 11, 2002

Date Due: Thursday, April 18, 2002 ****** Changed to Tuesday, April 23, 2002 ******

Reading: Please read Chapter 4, Sections 4.1 - 4.5, for April 18 and the week of April 22.

1. In this problem, I would like you to perform an analysis of a digital communication system that is similar to the case that we studied in class on April 16. The only difference is that the signal S and noise N have PMFs as follows.

$$P_S(s) = \begin{cases} 0.25, s = -1\\ 0.75, s = +1\\ 0, \text{ otherwise} \end{cases} \qquad P_N(n) = \begin{cases} 0.5, n = -1\\ 0.25, n = 0\\ 0.25, n = +1\\ 0, \text{ otherwise} \end{cases}$$

The random variables S and N are independent. The receiver observes the random variable X = S + N.

- (a) Suppose the receiver obtains the value X = x. Find the signal estimate $\hat{s}(x)$ that minimizes the *conditional* mean-squared error, $E[(S \hat{s}(x))^2 | X = x]$. Show all of the steps in your analysis, and display $\hat{s}(x)$ as a plot versus x. I suggest that you draw plots of $P_S(s)$, $P_N(n)$, $P_{X,S}(x,s)$, $P_X(x)$, $P_{S|X}(s|x)$ as we did in class.
- (b) How would you process the signal estimates $\hat{s}(x)$ in order to recover the binary values $\{-1, +1\}$ of S? In other words, what "decision rule" would you use to recover the bits from X based on $\hat{s}(x)$?
- (c) For the decision rule that you developed in part b, what is the probability of a bit error for this system? Explain your reasoning.
- (d) Write a MATLAB program to simulate this system, including the decision rule developed in part b. Compare the bit error rate (BER) in your simulation with the analytical probability of a bit error computed in part c.

Be sure to compare the simulated and analytical BER!

- 2. Please answer the following questions for the joint PMF $P_{X,Y}(x, y)$ shown in the figure on page 3.
 - (a) Find the marginal PMFs $P_X(x)$ and $P_Y(y)$, and plot them.
 - (b) Find the mean and variance of X and Y: $\mu_X, \mu_Y, \sigma_X^2, \sigma_Y^2$.
 - (c) Find the correlation between X and Y, defined as $r_{X,Y} = E[XY]$.
 - (d) Find the covariance Cov[X, Y].

- (e) Find the correlation coefficient $\rho_{X,Y}$.
- (f) Are X and Y independent random variables? Recall that X and Y are independent if and only if $P_{X,Y}(x,y) = P_X(x)P_Y(y)$.
- (g) Find the conditional PMFs

$$P_{X|Y}(x|y) = P[X = x|Y = y]$$
$$P_{Y|X}(y|x) = P[Y = y|X = x]$$

and plot each of these versus x and y (on separate plots).

- (h) Suppose you need to produce an estimate \hat{x} of X that minimizes the mean squared error $E[(X \hat{x})^2]$. You must do this with no information about Y. What value should you choose for \hat{x} , and why? (Your answer should be a number!)
- (i) Suppose that you *observe* that the random variable Y takes on the value y (i.e., Y = y). We would like to incorporate the knowledge that Y = y to improve our estimate of X. For each possible value of y, what is your estimate of X, denoted by $\hat{x}(y)$? Explain how to compute $\hat{x}(y)$, and present a plot of $\hat{x}(y)$ versus y.
- (j) Is $\hat{x}(y)$ in part (i) different from \hat{x} in part (h)? Is this reasonable based on the probability values in the joint PMF? Is this reasonable based on the value of correlation coefficient $\rho_{X,Y}$ that you computed in part (e)? That is, does the value of $\rho_{X,Y}$ lead you to expect that information about Y should be useful in predicting the value of X?
- (k) Using your answer from part (h), compute E[(X − x̂)²]. Using your answer from part (i), compute E[(X − x̂(−1))²|Y = −1], E[(X − x̂(0))²|Y = 0], and E[(X − x̂(1))²|Y = 1]. Do these results show that we get better estimates for X when the value of Y is known? Please explain.

